

TABLE 32. Sex Ratio of Fall-run Chinook Salmon and Coho Salmon Recovered During Carcass Surveys, South Fork Trinity River Basin, 1984 through 1990 (nd = no data).

Year	Number of carcasses recovered	Numbers (%)				Sex ratio (m:f)
		Males	Females	Sex not determined		
Fall-run Chinook Salmon						
1984	nd	nd	nd	nd	nd	nd
1985	160	124 (77.5%)	35 (21.9%)	1 (0.6%)		3.54
1986	184	118 (64.1%)	66 (35.9%)	0 (0.0%)		1.79
1987	76	53 (69.7%)	22 (28.9%)	1 (1.3%)		2.41
1988	49	38 (77.6%)	9 (18.4%)	2 (4.1%)		4.22
1989	77	43 (55.8%)	34 (44.2%)	0 (0.0%)		1.26
1990	74	64 (86.5%)	8 (10.8%)	2 (2.7%)		8.00
	Mean	71.9%	26.7%	1.5%		3.54
Coho Salmon						
1984	nd	nd	nd	nd	nd	nd
1985	35	6 (17.1%)	29 (82.9%)	0 (0.0%)		0.21
1986	1	1	0	0		
1987	3	2 (66.7%)	1 (33.3%)	0 (0.0%)		2.00
1988	0	0	0	0		
1989	9	4 (44.4%)	5 (55.6%)	0 (0.0%)		0.80
1990	20	8 (40.0%)	11 (55.0%)	1 (5.0%)		0.73
	Mean	42.1%	56.7%	1.3%		0.93